

f

act:

There's more than one way to fight type 2 diabetes.

f
un



f
ootwork




farxiga
(dapagliflozin)^{5mg} tablets

f
riends



f
lavor



See inside
for savings

Get the facts about food, fitness, and
THE PILL THAT STARTS WITH

f

FARXIGA® (dapagliflozin) is a prescription medicine used along with diet and exercise to lower blood sugar in adults with type 2 diabetes. FARXIGA should not be used to treat people with type 1 diabetes or diabetic ketoacidosis (increased ketones in your blood or urine).

Please see Important Safety Information throughout and on pages 6 and 7, and visit www.FARXIGAMedGuide.com for Medication Guide, and www.FARXIGAPI.com for US Full Prescribing Information for FARXIGA.

f

act:

FARXIGA may help you fight type 2 diabetes if you're not getting to goal on metformin

Along with diet and exercise, FARXIGA® (dapagliflozin), alone or in combination with other diabetes medicines, has been proven to help lower A1C in adults with type 2 diabetes.

If you're not reaching your A1C goal with metformin, diet, and exercise, talk to your doctor about adding FARXIGA, the pill that starts with f.

Ask your doctor about once-a-day FARXIGA.

For more information, visit FARXIGA.com

Important Safety Information for FARXIGA® (dapagliflozin)

Do not take FARXIGA if you:

- are allergic to dapagliflozin or any of the ingredients in FARXIGA. Symptoms of a serious allergic reaction may include skin rash, raised red patches on your skin (hives), swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing. If you have any of these symptoms, stop taking FARXIGA and contact your healthcare provider or go to the nearest hospital emergency room right away
- have severe kidney problems or are on dialysis. Your healthcare provider should do blood tests to check


farxiga®
(dapagliflozin) 5mg tablets



Important Safety Information for FARXIGA® (dapagliflozin)

how well your kidneys are working before and during your treatment with FARXIGA

FARXIGA may cause serious side effects including:

- Low blood sugar (hypoglycemia) can occur if you take FARXIGA with another medicine that can cause low blood sugar, such as sulfonylureas or insulin. Symptoms of low blood sugar include shaking, sweating, fast heartbeat, dizziness, hunger, headache, and irritability. Follow your healthcare provider's instructions for treating low blood sugar

Please see Important Safety Information on pages 6 and 7.

fact:

FARXIGA helps lower A1C in addition to diet and exercise

In clinical studies, FARXIGA:



Helped lower A1C*



Removed some blood sugar†

Additionally, FARXIGA:



Reduced weight on average 3%‡



Lowered systolic blood pressure‡

*Primary endpoint. †With FARXIGA at 12 weeks. ‡When used with metformin. Individual results may vary.

FARXIGA is not a weight-loss or blood-pressure drug.

#1
PRESCRIBED
WORLDWIDE

More than
30
CLINICAL
STUDIES

More than
1.8
MILLION
PATIENTS TREATED
WITH FARXIGA—

THE PILL THAT STARTS WITH **f**

FARXIGA is the #1 prescribed medicine of its kind worldwide, with more than 1.8 million patients currently on treatment. The benefits and safety of FARXIGA have been tested in more than 30 clinical studies of adults with type 2 diabetes.

FARXIGA, combined with diet and exercise, has been studied alone as well as in combination with other diabetes medicines you may be taking. The other medicines included metformin, glimepiride, pioglitazone, insulin, and sitagliptin.

Important Safety Information for FARXIGA® (dapagliflozin)

FARXIGA may cause serious side effects including:

- **Dehydration (the loss of body water and salt)**, which may cause you to feel dizzy, faint, lightheaded, or weak, especially when you stand up (orthostatic hypotension). You may be at a higher risk of dehydration if you have low blood pressure; take medicines to lower your blood pressure, including water pills (diuretics); are 65 years of age or older; are on a low salt diet, or have kidney problems

Important Safety Information for FARXIGA® (dapagliflozin)

FARXIGA may cause serious side effects including:

- **Low blood sugar (hypoglycemia)** can occur if you take FARXIGA with another medicine that can cause low blood sugar, such as sulfonylureas or insulin. Symptoms of low blood sugar include shaking, sweating, fast heartbeat, dizziness, hunger, headache, and irritability. Follow your healthcare provider's instructions for treating low blood sugar

fact:
FARXIGA works differently to rid your body of some sugar

FARXIGA removed approximately **70 grams of sugar** in the urine per day.*


*In a study of adults with type 2 diabetes at week 12 with FARXIGA.




How FARXIGA works

FARXIGA® (dapagliflozin) is part of a class of type 2 diabetes medications called SGLT2 (sodium-glucose cotransporter 2) inhibitors. It's just one pill a day taken in the morning with or without food.

Kidneys



1 Blocks some sugar from being absorbed by your body



2 Removes some sugar when your body has more than it needs



3 Flushes some sugar out of your body through your urine

Important Safety Information for FARXIGA® (dapagliflozin)

Do not take FARXIGA if you:

- have severe kidney problems or are on dialysis. Your healthcare provider should do blood tests to check how well your kidneys are working before and during your treatment with FARXIGA

Please see Important Safety Information throughout.

fact:

Most FARXIGA patients pay as low as \$0* a month

Savings and support: it's simple to register


Register for savings and support for FARXIGA. Give us a call at **1-855-3FARXIGA** or visit us at www.FARXIGAsavings.com



\$0* CO-PAY EVERY MONTH
for most patients[†]

Eligible commercially insured patients

BIN# 004682
PCN# CN
GRP# EC00000000
ID# XXXXXXXXXXXX



*As low as \$0 for as long as your doctor prescribes any available dose of FARXIGA.
†Data on File, US-8797, AstraZeneca Pharmaceuticals LP.

Please see accompanying US Full Prescribing Information, and Medication Guide.
Subject to eligibility. Restrictions apply. Not available for government-insured patients. See back of card for details.

Program managed by PSKW, LLC, on behalf of AstraZeneca. Fingertip Formulary is a registered trademark of Fingertip Formulary, LLC.

ELIGIBILITY: You may be eligible for this offer if you are insured by commercial insurance and your insurance does not cover the full cost of your prescription, or you are not insured and are responsible for the cost of your prescriptions. Patients who are enrolled in a state or federally funded prescription insurance program are not eligible for this offer. This includes patients enrolled in Medicare Part D, Medicaid, Medigap, Veterans Affairs (VA), Department of Defense (DOD) programs or TRICARE, and patients who are Medicare eligible and enrolled in an employer-sponsored group waiver health plan or government-subsidized prescription drug benefit program for retirees. If you are enrolled in a state or federally funded prescription insurance program, you may not use this savings card even if you elect to be processed as an uninsured (cash-paying) patient. This offer is not insurance, is restricted to residents of the United States and Puerto Rico, and to patients over 18 years of age.

TERMS OF USE: Eligible commercially insured patients with a valid prescription for FARXIGA® (dapagliflozin) who present this savings card at participating pharmacies will pay as low as \$0 per 30-day supply maximum savings limit applies; patient out-of-pocket expense may vary. If you pay cash for your prescription, AstraZeneca will pay up to the first \$150, and you will be responsible for any remaining balance, for each monthly prescription. Other restrictions may apply. Patient is responsible for applicable taxes, if any. Non-transferable, limited to one per person, cannot be combined with any other offer. Void where prohibited by law, taxed or restricted. Patients, pharmacists, and prescribers cannot seek reimbursement from health insurance or any third party for any part of the benefit received by the patient through this offer. AstraZeneca reserves the right to rescind, revoke, or amend this offer, eligibility and terms of use at any time without notice. This offer is not conditioned on any past, present or future purchase, including refills. Offer must be presented along with a valid prescription at the time of purchase. For additional details about this offer, please visit www.FARXIGAsavings.com. If you have any questions regarding this offer, please call **1-844-631-3978**.

BY USING THIS CARD, YOU AND YOUR PHARMACIST UNDERSTAND AND AGREE TO COMPLY WITH THESE ELIGIBILITY REQUIREMENTS AND TERMS OF USE.



fact:

Personalized support can help you stay motivated

Managing type 2 diabetes can be easier when you get help along the way. When you sign up for support from FARXIGA® (dapagliflozin), you'll receive a welcome package and a series of e-mails with:

- Resources to help you move toward your goals
- Tools with insights about diet, exercise, and more
- Information about living with type 2 diabetes

Sign up today at www.FARXIGAsavings.com

FARXIGA Care Partners* Get connected to a variety of support services, tips, and information with the help of FARXIGA Care Partners.

FARXIGA Prescription Coverage Counselors†

We have Prescription Coverage Counselors who can talk to you on the phone about your FARXIGA coverage and benefit verification. They can help you determine your co-pay amount and what information you'll need to process your benefit verification.

Visit [FARXIGA.com](http://www.FARXIGA.com)

or call

1-855-3FARXIGA
to find out more.

*FARXIGA Care Partners cannot provide medical advice and can only provide information about AstraZeneca medications. Decisions regarding your health and treatment of your condition should be made with your own healthcare provider.

†Prescription Coverage Counselors do not file claims or appeals on behalf of patients or healthcare professionals. Coverage or co-pay amounts may change prior to the time of actual prescription fill.

Important Safety Information for FARXIGA® (dapagliflozin)

Who should not take FARXIGA?

Do not take FARXIGA if you:

- are allergic to dapagliflozin or any of the ingredients in FARXIGA. Symptoms of a serious allergic reaction may include skin rash, raised red patches on your skin (hives), swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing. If you have any of these symptoms, stop taking FARXIGA and contact your healthcare provider or go to the nearest hospital emergency room right away
- have severe kidney problems or are on dialysis. Your healthcare provider should do blood tests to check how well your kidneys are working before and during your treatment with FARXIGA

What are the possible side effects of FARXIGA?

FARXIGA may cause serious side effects including:

- **Dehydration** (the loss of body water and salt), which may cause you to feel dizzy, faint, lightheaded, or weak, especially when you stand up (orthostatic hypotension). You may be at a higher risk of dehydration if you have low blood pressure; take medicines to lower your blood pressure, including water pills (diuretics); are 65 years of age or older; are on a low salt diet, or have kidney problems
- **Ketoacidosis** occurred in people with type 1 and type 2 diabetes during treatment with FARXIGA. Ketoacidosis is a serious condition which may require hospitalization and may lead to death. Symptoms may include nausea, tiredness, vomiting, trouble breathing, and abdominal pain. If you get any of these symptoms, stop taking FARXIGA and call your healthcare provider right away. If possible, check for ketones in your urine or blood, even if your blood sugar is less than 250 mg/dL

Please visit www.FARXIGAMedGuide.com for Medication Guide, and www.FARXIGAPI.com for US Full Prescribing Information for FARXIGA.

Important Safety Information for FARXIGA® (dapagliflozin) (cont'd)

- **Kidney problems.** Sudden kidney injury occurred in people taking FARXIGA. Talk to your doctor right away if you reduce the amount you eat or drink, or if you lose liquids; for example, from vomiting, diarrhea, or excessive heat exposure
- **Serious urinary tract infections (UTI),** some that lead to hospitalization, occurred in people taking FARXIGA. Tell your doctor if you have any signs or symptoms of UTI including a burning feeling when passing urine, a need to urinate often, the need to urinate right away, pain in the lower part of your stomach (pelvis), or blood in the urine with or without fever, back pain, nausea, or vomiting
- **Low blood sugar (hypoglycemia)** can occur if you take FARXIGA with another medicine that can cause low blood sugar, such as sulfonylureas or insulin. Symptoms of low blood sugar include shaking, sweating, fast heartbeat, dizziness, hunger, headache, and irritability. Follow your healthcare provider's instructions for treating low blood sugar
- **Vaginal yeast infections** in women who take FARXIGA. Talk to your healthcare provider if you experience vaginal odor, white or yellowish vaginal discharge (discharge may be lumpy or look like cottage cheese), or vaginal itching
- **Yeast infection of skin around the penis (balanitis)** in men who take FARXIGA. Talk to your healthcare provider if you experience redness, itching, or swelling of the penis; rash of the penis; foul smelling discharge from the penis; or pain in the skin around penis. Certain uncircumcised men may have swelling of the penis that makes it difficult to pull back the skin around the tip of the penis
- **Increase in bad cholesterol (LDL-C).** Your healthcare provider should check your LDL-C during treatment with FARXIGA
- **Bladder cancer.** In studies of FARXIGA in people with diabetes, bladder cancer occurred in a few more people who were taking FARXIGA than in people who were taking other diabetes medication. There were too few cases of bladder cancer to know if bladder cancer was related to FARXIGA. Tell your healthcare provider right away if you have blood or a red color in your urine or pain while you urinate

Important Safety Information for FARXIGA® (dapagliflozin) (cont'd)

The most common side effects of FARXIGA include yeast infections of the vagina or penis, and changes in urination, including urgent need to urinate more often, in larger amounts, or at night.

What should I tell my healthcare provider before taking FARXIGA?

Before you take FARXIGA, tell your healthcare provider:

- **all of your medical conditions,** including problems with your kidneys, liver, bladder, or pancreas
- **if you have had, or have risk factors for, ketoacidosis** (including type 1 diabetes, are eating less due to illness, surgery, or a change in your diet, are going to have surgery, or binge drink)
- **if you are pregnant, or plan to become pregnant.** FARXIGA may harm your unborn baby
- **if you are breastfeeding, or plan to breastfeed.** It is unknown if FARXIGA passes into your breast milk
- **about all the medicines you take,** including prescription and nonprescription medicines, vitamins, and herbal supplements

Approved Uses for FARXIGA® (dapagliflozin)

What is FARXIGA?

FARXIGA is a prescription medicine used along with diet and exercise to lower blood sugar in adults with type 2 diabetes.

FARXIGA should not be used to treat people with type 1 diabetes or diabetic ketoacidosis (increased ketones in your blood or urine).

Please visit www.FARXIGAMedGuide.com for Medication Guide, and www.FARXIGAPI.com for US Full Prescribing Information for FARXIGA.

You may report side effects related to AstraZeneca products by clicking [here](#).

If you can't afford your medication, AstraZeneca may be able to help. For more information, please visit www.AstraZeneca-us.com.

f

act:

Not everyone can control type 2 diabetes with food, fitness, and metformin

Questions to ask your doctor if you haven't reached your goals

About your A1C

1. What is my A1C now?
2. What is my A1C goal?
3. How long should it take me to meet my A1C goal?

About FARXIGA® (dapagliflozin)

1. Could including FARXIGA help me meet my goal?
2. How does FARXIGA work to reduce blood sugar?
3. In what other ways could FARXIGA help me?

Things to ask at the end of your appointment

1. What changes were made to my treatment plan?
2. How often should I check my blood sugar?
3. When should I schedule my next appointment?

Hot down your thoughts to share with your doctor


farxiga[®]
(dapagliflozin) ^{5mg}
tablets



Ask your
doctor about

THE PILL THAT STARTS WITH **f**

Visit [FARXIGA.com](https://www.farxiga.com) to learn how you can
save on your prescription.

AstraZeneca 

FARXIGA is a registered trademark of the
AstraZeneca group of companies.

©2018 AstraZeneca. All rights reserved.
US-18869 Last Updated 4/18